

VOLTANEC SP Tablets (Aceclofenac + Paracetamol + Serratiopeptidase)

Black Box Warning

Hepatotoxicity

Taking more than daily dose of Paracetamol may cause serious liver damage or allergic reactions (eg. swelling of the face, mouth and throat, difficulty in breathing, itching or rash). Most of the cases of liver injury are associated with the use of paracetamol at doses that exceed 4,000 mg per day. (See WARNINGS AND PRECAUTIONS)

Composition

VOLTANEC-SP Tablets

Each film-coated tablet contains:

Aceclofenac, IP100 mg

Paracetamol, IP 325 mg

Serratiopeptidase, IP15 mg

eg. to enzyme activity of 30,000 units

(as enteric-coated granules)

Dosage Form

Tablet for oral use

Description

VOLTANEC-SP is a fixed-dose combination of aceclofenac 100 mg, paracetamol 500 mg and serratiopeptidase 15 mg. This combination offers the advantage of the peripheral effects of aceclofenac, the central effect of paracetamol and pain relief with the tissue repair effects of serratiopeptidase for pain management.

Aceclofenac is an orally administered phenylacetic acid derivative with effects on a variety of inflammatory mediators. Through its analgesic and anti-inflammatory properties, aceclofenac provides symptomatic relief in a variety of painful conditions. Due to its preferential cyclooxygenase (COX-2) blockade, it has better safety than conventional non-steroidal anti-inflammatory drugs (NSAIDs) with respect to adverse effects on the gastrointestinal and cardiovascular system.

Paracetamol acts predominantly by inhibiting prostaglandin synthesis in the central nervous system and, to a lesser extent, through a peripheral action by blocking pain impulse generation. Paracetamol is one of the effective mild analgesics, suitable for treating mild-to-moderate pain.

Serratiopeptidase breaks down fibrin, which is believed to be one of the causes of chronic inflammation. It also causes thinning of the fluids in the body as a result of injury, making fluid drainage smoother. It can

break down protein and facilitate fibrinolytic activity in the blood, and modify molecules that guide inflammatory cells to their targets. This activity includes the digestion of scar tissue, blood clots, cysts, mucus, arterial plaque, and all types of inflammation. Serratiopeptidase reduces internal tissue oedema and inflammation caused at post-operating handling. It can also block the pain-inducing amines from swollen tissues.

Pharmacology

► Pharmacodynamics

Aceclofenac

Aceclofenac is a non-steroidal agent with marked anti-inflammatory and analgesic properties. The mode of action of aceclofenac is largely based on the inhibition to prostaglandin synthesis. Aceclofenac is a potent inhibitor of the enzyme, COX, which is involved in the production of prostaglandins.

Aceclofenac relieves pain and inflammation through a variety of mechanisms and, in addition, exerts stimulatory effects on cartilage matrix synthesis.

Anti-inflammatory Activity: The anti-inflammatory effects of aceclofenac have been shown in both acute and chronic inflammation. It inhibits various mediators of pain and inflammation, including the following:

- *PGE2 via COX inhibition* (COX-1 and COX-2) after intracellular metabolism to 4-hydroxyaceclofenac and diclofenac in human rheumatoid synovial cells and other inflammatory cells.
- *IL-1beta, IL-6 and tumour necrosis factor-alpha* in human osteoarthritic synovial cells and human articular chondrocytes.
- *Reactive oxygen species* (which plays a role in joint damage) has also been observed in patients with osteoarthritis of the knees.
- *Expression of cell adhesion molecules* (which is implicated in cell migration and inflammation) has also been shown in human neutrophils.

Stimulatory Effects on Cartilage Matrix Synthesis: Aceclofenac stimulates glycosaminoglycan synthesis in human osteoarthritic cartilage by inhibition of IL-1beta and suppresses cartilage degeneration by inhibiting IL-1beta-mediated promatrix metalloproteinase production and proteoglycan release.

Paracetamol

Paracetamol is an aniline derivative with analgesic and antipyretic actions similar to those of aspirin, but with no demonstrable anti-inflammatory activity. Paracetamol is less irritant to the stomach than aspirin. It does not affect thrombocyte aggregation or bleeding time. Paracetamol is generally well tolerated by patients hypersensitive to acetylsalicylic acid.

Analgesic Action: The central analgesic action of paracetamol resembles that of aspirin. It produces analgesia by raising the pain threshold.

Antipyretic Effect: The antipyretic effect of paracetamol is attributed to its ability to inhibit COX in the brain where the peroxide tone is low. Recent evidence suggests inhibition of COX-3 (believed to be a splice variant product of the COX-1 gene) and could represent a primary central mechanism by which paracetamol decreases pain and, possibly, fever.

Serratiopeptidase

It binds to alpha-2-macroglobulin in the blood in the ratio of 1:1, which helps to mask its antigenicity, but retain its enzymatic activity. Levels of serratiopeptidase are slowly transferred to the exudate at the site of inflammation and gradually, the blood level declines. By hydrolysing bradykinin, histamine and serotonin, it indirectly reduces dilatation of blood capillaries and controls permeability. Serratiopeptidase blocks plasmin inhibitors, thus helping the fibrinolytic activity of plasmin. Degradation of extra-fibrin to small fragment

prevents the clogging of microcapillaries, helps clearance of exudates, reduces swelling and improves microcirculation.

► Pharmacokinetics

Aceclofenac

Absorption

After oral administration, aceclofenac is rapidly and completely absorbed as unchanged drug. Peak plasma concentrations are reached approximately 1.25-3.00 hours following ingestion.

Distribution

Aceclofenac penetrates into the synovial fluid, where the concentrations reach approximately 57% of those in plasma. The volume of distribution is approximately 25 L. Aceclofenac is highly protein-bound (>99%). Aceclofenac circulates mainly as unchanged drug.

Metabolism

4-hydroxyaceclofenac is the main metabolite detected in plasma.

Elimination

The mean plasma elimination half-life is around 4 hours. Approximately two-thirds of the administered dose is excreted via the urine, mainly as hydroxymetabolites.

Paracetamol

Absorption

Paracetamol is well absorbed by the oral route. The plasma half-life is about 2 hours.

Distribution

Plasma protein binding is negligible at the usual therapeutic concentration, but increases with increasing concentrations. Acetaminophen is, relatively, uniformly distributed throughout most body fluids. The plasma half-life is ($t_{1/2}$) 2-3 hours and the effect after an oral dose lasts for 3-5 hours.

Metabolism

Paracetamol is primarily metabolized in the liver by conjugation to glucuronide and sulphate. A small amount (about 3-10% of a therapeutic dose) is metabolized by oxidation and the reactive intermediate metabolite thus formed is bound preferentially to the liver glutathione and excreted as cysteine and mercapturic acid conjugates.

Elimination

Excretion occurs via the kidneys. Of a therapeutic dose, 2-3% is excreted unchanged, 80-90% as glucuronide and sulphate, and a smaller amount as cysteine and mercapturic acid derivatives.

Serratiopeptidase

Absorption

After oral administration, serratiopeptidase is almost totally absorbed from the gastrointestinal (GI) tract.

Distribution

Serratiopeptidase binds to alpha-2 macroglobulin in the blood and produces an enzyme activity in the blood circulation. It shows a steep rise in concentration at the site of injury and inflammation.

Metabolism

Metabolism of serratiopeptidase takes place in the liver.

Excretion

The metabolites of serratiopeptidase are excreted through the urine and faeces.

Indications

- Resolution of inflammation and pain due to bone and soft tissue injury.

- Resolution of post-operative inflammation, oedema and pain.

Dosage And Administration

VOLTANEC-SP is supplied for oral administration in adults and should be swallowed whole with a sufficient amount of liquid. It should be taken preferably with or after food.

The maximum recommended dose of VOLTANEC-SP is two tablets daily, taken as one tablet in the morning and one in the evening.

Contraindications

VOLTANEC-SP is contraindicated in the following situations:

- Patients sensitive to aceclofenac, paracetamol, serratiopeptidase or to any of the excipients of the product.
- Patients with a history of or active, recurrent peptic ulcer/haemorrhage (two or more distinct episodes of proven ulceration or bleeding).
- Patients who have previously shown hypersensitivity reactions (e.g. asthma, rhinitis, angio-oedema or urticaria) in response to ibuprofen, aspirin or other NSAIDs.
- Patients with a history of anaphylactic reactions.
- Patients with severe heart failure, hypertension, and hepatic or renal impairment should not be prescribed.
- During pregnancy, especially during the last trimester of pregnancy, unless there are compelling reasons for doing so. The lowest effective dosage should be used.

Warnings And Precautions

► General

Undesirable effects may be minimized by using the lowest effective dose for the shortest duration necessary to control symptoms. Concomitant use with NSAIDs, including COX-2 selective inhibitors, should be avoided. It should not be combined with other analgesic medications that contain paracetamol and should be given with care to patients with impaired kidney or liver function.

The administration of an NSAID may cause a dose-dependent reduction in prostaglandin formation and precipitate renal failure. Patients at the greatest risk of this reaction are those with impaired renal function, cardiac impairment, liver dysfunction, those taking diuretics and the elderly. Renal function should be monitored in these patients.

Respiratory Disorders

Caution is required if administered to patients suffering from, or with a previous history of, bronchial asthma since NSAIDs have been reported to precipitate bronchospasm in such patients.

Hepatic Toxicity (Also SEE BLACK BOX WARNING)

Paracetamol may cause liver damage if more than the recommended dose is taken. Allergic reactions like swelling of the face, mouth and throat, difficulty in breathing, itching or rash may occur due to high doses of paracetamol. Severe liver damage may occur if:

- Adult takes more than 4000 mg in 24 hours, which is the maximum daily amount
- Child takes more than 5 doses in 24 hours
- Taken with other drugs containing paracetamol
- Adult has 3 or more alcoholic drinks every day while using this product

Cardiovascular and Cerebrovascular Effects

Appropriate monitoring and advice are required for patients with a history of hypertension and/or mild-t-

-moderate congestive heart failure as fluid retention and oedema have been reported in association with NSAID therapy.

Clinical trial and epidemiological data suggest that the use of some NSAIDs (particularly at high doses and in long-term treatment) may be associated with a small increased risk of arterial thrombotic events (e.g., myocardial infarction or stroke). There are insufficient data to exclude such a risk for aceclofenac.

Patients with uncontrolled hypertension, congestive heart failure, established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease should only be treated after careful consideration. Similar consideration should be made before initiating long-term treatment of patients with risk factors for cardiovascular disease (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking).

GI Bleeding, Ulceration and Perforation

GI bleeding, ulceration or perforation, which can be fatal, has been reported with all NSAIDs at any time during treatment, with or without warning symptoms or a previous history of serious GI events.

Close medical surveillance is imperative in patients with symptoms indicative of GI disorders, with a history suggestive of GI ulceration, with ulcerative colitis or with Crohn's disease, bleeding diathesis or haematological abnormalities.

The risk of GI bleeding, ulceration or perforation is higher with increasing NSAID doses, in patients with a history of ulcers, particularly if complicated with haemorrhage or perforation, and in the elderly. These patients should commence treatment on the lowest dose available. Combination therapy with protective agents (e.g. misoprostol or proton-pump inhibitors) should be considered for these patients, and also for patients requiring concomitant low-dose aspirin, or other drugs likely to increase GI risk.

Patients with a history of GI toxicity, particularly when elderly, should report any unusual abdominal symptoms (especially GI bleeding), particularly in the initial stages of treatment.

Caution should be advised in patients receiving concomitant medications that could increase the risk of ulceration or bleeding, such as oral corticosteroids, anticoagulants such as warfarin, selective serotonin-reuptake inhibitors or antiplatelet agents such as aspirin. When GI bleeding or ulceration occurs, the treatment should be withdrawn.

NSAIDs should be given with care to patients with a history of GI disease (ulcerative colitis, Crohn's disease) as these conditions may be exacerbated.

SLE and Mixed Connective Tissue Disorders

In patients with systemic lupus erythematosus (SLE) and mixed connective tissue disorders, there may be an increased risk of aseptic meningitis.

Dermatological

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs. Patients appear to be at the highest risk for these reactions early in the course of therapy, with the onset of the reaction occurring in the majority of cases within the first month of treatment. Discontinuation should be done at the first appearance of skin rash, mucosal lesions or any other sign of hypersensitivity.

Hypersensitivity Reactions

As with other NSAIDs, allergic reactions, including anaphylactic/anaphylactoid reactions, can also occur without earlier exposure to the drug.

Haematological

May cause reversibly inhibit platelet aggregation.

Long-term Treatment

Individuals receiving long-term treatment should be regularly monitored for renal function tests, liver function tests and blood counts.

It is to be used with caution in hepatic porphyria, coagulation disorders, history of peptic ulcers, ulcerative colitis, Crohn's disease, cerebrovascular bleeding, pregnancy and lactation. Caution should be exercised in patients with mild-to-moderate impairment of cardiac, hepatic or renal function and in elderly patients who are more likely to be suffering from these conditions. Caution is also required in patients on diuretic therapy or otherwise at risk of hypovolaemia.

It may cause dizziness. Driving or operating machinery is to be avoided.

▶ Drug Interactions

Aceclofenac

Other Analgesics, Including cox-2 Selective Inhibitors: Avoid concomitant use of two or more NSAIDs (including aspirin) as this may increase the risk of adverse effects.

Antihypertensives: Reduced antihypertensive effect.

Diuretics: Reduced diuretic effect. Diuretics can increase the risk of nephrotoxicity of NSAIDs. Although it was not shown to affect blood pressure control when co-administered with bendrofluazide, interactions with other diuretics cannot be ruled out. When concomitant administration with potassium-sparing diuretics is employed, serum potassium should be monitored.

Cardiac Glycosides: NSAIDs may exacerbate cardiac failure, reduce the glomerular filtration rate (GFR) and increase plasma glycoside levels.

Lithium: Decreased elimination of lithium.

Methotrexate: Decreased elimination of methotrexate. Caution should be exercised if NSAIDs and methotrexate are administered within 24 hours of each other, since NSAIDs may increase plasma levels, resulting in increased toxicity.

Ciclosporin: Increased risk of nephrotoxicity.

Mifepristone: NSAIDs should not be used for 8-12 days after mifepristone administration as NSAIDs can reduce the effect of mifepristone.

Corticosteroids: Increased risk of GI ulceration or bleeding.

Anticoagulants: NSAIDs may enhance the effects of anticoagulants, such as warfarin. Close monitoring of patients on combined anticoagulants and aceclofenac therapy should be undertaken.

Quinolone Antibiotics: Animal data indicate that NSAIDs can increase the risk of convulsions associated with quinolone antibiotics. Patients taking NSAIDs and quinolones may have an increased risk of developing convulsions. Antiplatelet agents and selective serotonin-reuptake inhibitors can lead to increased risk of GI bleeding.

Tacrolimus: Possible increased risk of nephrotoxicity when NSAIDs are given with tacrolimus.

Zidovudine: Increased risk of haematological toxicity when NSAIDs are given with zidovudine. There is evidence of an increased risk of haemarthroses and haematoma in HIV(+) haemophiliacs receiving concurrent treatment with zidovudine and ibuprofen.

Antidiabetic Agents: Clinical studies have shown that diclofenac can be given together with oral antidiabetic agents without influencing their clinical effect. However, there have been isolated reports of hypoglycaemic and hyperglycaemic effects. Thus, with aceclofenac, consideration should be given to adjustment of the dosage of hypoglycaemic agents.

Other NSAIDs: Concomitant therapy with aspirin or other NSAIDs may increase the frequency of adverse reactions, including the risk of GI bleeding.

Paracetamol

Drugs that induce hepatic microsomal enzymes, such as alcohol, barbiturates and other anticonvulsants, may increase the hepatotoxicity of paracetamol, particularly after overdose.

The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular use of paracetamol, with increased risk of bleeding. The effect appears to increase as the dose of paracetamol is increased, but can occur with doses as low as 1.5-2 g paracetamol per day for at least 5-7 days. Occasional doses have no significant effect.

Probenicid inhibits the glucuronidation of paracetamol which can affect the clearance of paracetamol. This should be considered when these medicines are administered concomitantly.

Paracetamol may affect the pharmacokinetics of chloramphenicol. This interaction should be considered when these medications are administered concomitantly, especially in malnourished patients.

Enzyme-inducing medicines, such as some antiepileptic drugs (phenytoin, phenobarbital, carbamazepine) have been shown in pharmacokinetic studies to reduce the plasma AUC of paracetamol to approximately 60%. Other substances with enzyme-inducing properties, e.g. rifampicin and St John's wort (*Hypericum perforatum*) are also suspected of causing lowered concentrations of paracetamol. In addition, the risk of liver damage during treatment with the maximum recommended doses of paracetamol will be higher in patients being treated with enzyme-inducing agents.

Serratiopeptidase

Serratiopeptidase may interact with medications that slow blood clotting (anticoagulant/antiplatelet drugs). Serratiopeptidase might decrease blood clotting. Therefore, taking serratiopeptidase along with medications that also slow clotting might increase the chances of bruising and bleeding. Some medications that slow blood clotting include aspirin, clopidogrel, diclofenac, ibuprofen, naproxen, dalteparin, enoxaparin, heparin, warfarin and others.

► Renal Impairment

It should be avoided in patients with moderate and severe renal impairment. The importance of prostaglandins in maintaining renal blood flow should be taken into account in patients with impaired cardiac or renal function, those being treated with diuretics or recovering from major surgery. Effects on renal function are usually reversible on withdrawal.

► Hepatic Impairment

In patients with hepatic impairment, dosage reductions are recommended. If abnormal liver function tests persist or worsen, clinical signs or symptoms consistent with liver disease develop or if other manifestations occur (eosinophilia, rash), it should be discontinued. Close medical surveillance is necessary in patients suffering from mild to moderate impairment of hepatic function. Hepatitis may occur without prodromal symptoms. Use in patients with hepatic porphyria may trigger an attack.

► Geriatric

As with other NSAIDs and combinations, caution is advised in elderly patients who are more likely to have concomitant renal, hepatic or cardiovascular impairment or receiving concurrent medication. The elderly have an increased frequency of adverse reactions to NSAIDs, especially GI bleeding and perforation, which may be fatal.

► Pregnancy

Congenital abnormalities have been reported in association with NSAID administration in humans; however, these are low in frequency and do not appear to follow any discernible pattern. In view of the known effects of NSAIDs on the foetal cardiovascular system (risk of closure of the ductus arteriosus) and on the possible risk of persistent pulmonary hypertension of the newborn, use in the last trimester of pregnancy is contraindicated. The regular use of NSAIDs during the last trimester of pregnancy may decrease uterine tone

and contraction. The onset of labour may be delayed and the duration increased, with an increased bleeding tendency in both mother and child. NSAIDs should not be used during the first two trimesters of pregnancy or labour unless the potential benefit to the patient outweighs the potential risk to the foetus. The drug is not recommended in pregnant women.

▶ Lactation

In limited studies so far available, NSAIDs can appear in breast milk in very low concentrations. NSAIDs should, if possible, be avoided when breastfeeding.

The use of aceclofenac should be avoided in pregnancy and lactation unless the potential benefits to the other outweigh the possible risks to the foetus. The drug is not recommended in breastfeeding women.

Undesirable Effects

▶ GI Effects

The most commonly-observed adverse events are GI in nature. Peptic ulcers, perforation or GI bleeding, sometimes fatal, particularly in the elderly, may occur. Nausea, vomiting, diarrhoea, flatulence, constipation, dyspepsia, abdominal pain, melaena, haematemesis, ulcerative stomatitis, exacerbation of colitis and Crohn's disease have been reported following administration. Less frequently, gastritis has been observed. Pancreatitis has been reported very rarely.

▶ Hypersensitivity

Hypersensitivity reactions have been reported following treatment with NSAIDs. These may consist of (a) non-specific allergic reactions and anaphylaxis (b) respiratory tract reactivity comprising asthma, aggravated asthma, bronchospasm or dyspnoea, or (c) assorted skin disorders, including rashes of various types, pruritus, urticaria, purpura, angio-oedema and, more rarely, exfoliative and bullous dermatoses (including epidermal necrolysis and erythema multiforme).

▶ Cardiovascular

Oedema, hypertension and cardiac failure have been reported in association with NSAID treatment. Clinical trial and epidemiological data suggest that use of some NSAIDs (particularly at high doses and in long-term treatment) may be associated with an increased risk of arterial thrombotic events (e.g. myocardial infarction or stroke). Vasculitis has been reported rarely.

▶ Other Adverse Reactions Reported Less Commonly

Renal: Nephrotoxicity in various forms, including interstitial nephritis, nephritic syndrome and renal failure.

Hepatic: Abnormal liver function, hepatitis and jaundice.

Neurological and Special Senses: Visual disturbances, optic neuritis, headaches, paraesthesia, reports of aseptic meningitis (especially in patients with existing autoimmune disorders, such as SLE and mixed connective tissue disorders), with symptoms such as stiff neck, headache, nausea, vomiting, fever or disorientation, depression, confusion, hallucinations, tinnitus, vertigo, dizziness, malaise, fatigue and drowsiness.

Haematological: Thrombocytopenia, neutropenia, agranulocytosis, aplastic anaemia and haemolytic anaemia.

Dermatological: Bullous reactions, including Stevens-Johnson syndrome and toxic epidermal necrolysis (very rare). Photosensitivity.

Within the system organ classes, undesirable effects are listed under headings of frequency, using the

following categories: very common (1/10); common (1/100 to <1/10); uncommon (1/1,000 to <1/100); rare (1/10,000 to <1/1,000); very rare (<1/10,000), and not known (cannot be estimated from the available data). Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System organ class	Common ($\geq 1/100$ to <1/10)	Uncommon ($\geq 1/1,000$ to <1/100)	Rare ($\geq 1/10,000$ to <1/1,000)	Very rare/ isolated reports (<1/10,000)
Blood and lymphatic system disorders			Anaemia	Granulocytopenia Thrombocytopenia Neutropenia Haemolytic anaemia
Immune system disorders			Anaphylactic reaction (including shock) Hypersensitivity Allergic reaction	
Metabolism and nutrition disorders				Hyperkalemia Hypoglycaemia
Psychiatric disorders				Depression Abnormal dreams Insomnia
Nervous system disorders	Dizziness, drowsiness			Paraesthesia Tremor Somnolence Headache Dysgeusia (abnormal taste)
Eye disorders			Visual disturbance	
Ear and labyrinth disorders				Vertigo
Cardiac disorders				Palpitations
Vascular disorders				Flushing Hot flush
Respiratory, thoracic and mediastinal disorders			Dyspnoea	Bronchospasm Stridor

GI disorders	Dyspepsia Abdominal pain Nausea Diarrhoea Redness of the rectal mucous membranes	Flatulence Gastritis Constipation Vomiting Mouth ulceration	Melaena	Stomatitis Haematemesis GI haemorrhage Gastric ulcer Pancreatitis
Hepatobiliary disorders				Hepatitis Jaundice Liver damage
Skin and subcutaneous tissue disorders		Pruritus Rash Dermatitis Urticaria	Face oedema Exanthema Urticarial	Purpura Dermatitis bullous Exanthema Angioedema
Musculoskeletal and connective tissue disorders				Cramps in the leg
Renal and urinary disorders				Renal impairment Nephrotic syndrome
General disorders and administration site conditions				Oedema Fatigue Cramps in legs
Investigations	Hepatic enzyme increased	Blood urea increased Blood creatinine increased		Blood alkaline phosphatase increased Weight increase Increased serum potassium

Most of the adverse events are minor and reversible with treatment discontinuation. As with other NSAIDs, severe mucocutaneous skin reactions may also occur.

Overdosage

Liver damage is possible in adults who have taken 10 g or more of paracetamol. Ingestion of 5 g or more of paracetamol may lead to liver damage if the patient has risk factors.

► Risk factors

If the patient

- is on long term treatment with carbamazepine, phenobarbitone, phenytoin, primidone, rifampicin, St John's wort or other drugs that induce liver enzymes, or
- is likely to be glutathione depleted e.g. eating disorders, cystic fibrosis, HIV infection, starvation, cachexia.

Immediate treatment is essential in the management of overdosage. Despite a lack of significant early

symptoms, patients should be referred to a hospital urgently for immediate medical attention. Symptoms include headache, pallor, nausea, vomiting, epigastric pain, GI irritation, GI bleeding, rarely diarrhoea, disorientation, excitation, coma, drowsiness, dizziness, tinnitus, hypotension, respiratory depression, fainting, occasionally convulsions, anorexia and abdominal pain. In cases of significant poisoning, acute renal failure and liver damage are possible. Abnormalities of glucose metabolism and metabolic acidosis may occur. In severe poisoning, hepatic failure may progress to encephalopathy, coma and death. Acute renal failure with acute tubular necrosis may develop even in the absence of severe liver damage. Cardiac arrhythmias and pancreatitis have been reported.

Patients should be treated symptomatically as required. Within 1 hour of ingestion of a potentially toxic amount, activated charcoal should be considered. Alternatively, in adults, gastric lavage should be considered within 1 hour of ingestion of a potentially life-threatening overdose. Administration of oral methionine or intravenous N-acetylcysteine, which may have a beneficial effect up to at least 48 hours after the overdose, may be required. General supportive measures must be available.

Specific therapies such as dialysis or haemoperfusion are probably of no help in eliminating NSAIDs due to their high rate of protein binding and extensive metabolism. Good urine output should be ensured.

Renal and liver function should be closely monitored. Patients should be observed for at least 4 hours after ingestion of potentially toxic amounts. In case of frequent or prolonged convulsions, patients should be treated with intravenous diazepam.

Other measures may be indicated by the patient's clinical condition. Management of acute poisoning with NSAIDs essentially consists of supportive and symptomatic measures. Management of acute poisoning with NSAIDs essentially consists of supportive and symptomatic measures.

Storage And Handling Instructions

Store in a cool, dry place. Protect from light.

Packaging Information

VOLTANEC-SP is available in a strip pack of 10 tablets.

Last updated: November 2013

Last reviewed: November 2013

VOLTANEC SP Tablets

Source URL: <https://ciplamed.com/content/voltanec-sp-tablets>